

Triple Neutral Gauge Boson Couplings in Noncommutative Standard Model

¹N.G. Deshpande and ²Xiao-Gang He

¹ Institute of Theoretical Sciences, University of Oregon, Eugene, OR97403, USA

and

²Department of Physics, National Taiwan University, Taipei, Taiwan 10764, R.O.C.

Abstract

It has been shown recently that the triple neutral gauge boson couplings are not uniquely determined in noncommutative extension of the Standard Model (NCSM). Depending on specific schemes used, the couplings are different and may even be zero. To distinguish different realizations of the NCSM, additional information either from theoretical or experimental considerations is needed. In this paper we show that these couplings can be uniquely determined from considerations of unification of electroweak and strong interactions. Using $SU(5)$ as the underlying theory and integrating out the heavy degrees of freedom, we obtain unique non-zero new triple $\gamma\gamma\gamma$, $\gamma\gamma Z$, γZZ , ZZZ , γGG , ZGG and GGG couplings at the leading order in the NCSM. We also briefly discuss experimental implications.

PACS numbers:

Typeset using REVTeX

The property of space-time has fundamental importance in understanding the law of nature. Noncommutative (NC) quantum field theory, which modifies the space-time commutation relations, provides an alternative to the ordinary quantum field theory which may shed some light on the detailed structure of space-time. A simple way to modify the space-time properties is to change the usual space-time coordinate x to noncommutative coordinate $\hat{X}$ such that [1]

$$[\hat{X}^\mu, \hat{X}^\nu] = i\theta^{\mu\nu}. \quad (1)$$

We consider the case with $\theta^{\mu\nu}$ be a constant real anti-symmetric matrix which commutes with $\hat{X}^\mu$.

NC quantum field theory based on the above commutation relation can be easily studied using the Weyl-Moyal correspondence replacing the product of two fields $A(\hat{X})$ and $B(\hat{X})$ with NC coordinates by the star “*” product [2],

$$A(\hat{X})B(\hat{X}) \rightarrow A(x) * B(x) = \text{Exp}[i\frac{1}{2}\theta^{\mu\nu}\partial_{x,\mu}\partial_{y,\nu}]A(x)B(y)|_{x=y}. \quad (2)$$

Phenomenology for NC electromagnetic theory $U(1)_{em}$ has been studied recently [3], but less has been done for noncommutative Standard Model (NCSM) because it is non-trivial to construct such a theory. Due to the noncommuting nature of the “*” product, even with a $U(1)$ gauge theory the charges of matter fields in the theory are fixed to only three possible values, 1, 0, -1 [4]. Also $SU(N)$ group can not simply be gauged with “*” product, but $U(N)$ can be [4]. These pose certain difficulties in constructing NCSM, since in the SM the $U(1)_Y$ charges are not just 1, 0, -1, but some of them are fractionally charged, such as 1/6, 2/3, -1/3 for left handed quarks, right handed up and down quarks, respectively. There are also the problems with gauging the $SU(3)_C$ and $SU(2)_L$ groups. However, all these difficulties can be overcome with the use of the Seiberg-Witten map [5] which maps noncommutative $U(N)$ gauge fields to commutative $SU(N)$ gauge fields. Therefore, consistent noncommutative $SU(N)$ gauge theory can be constructed. The same map can also cure the charge quantization problem as shown in Ref. [6,7] by introducing new degrees of freedom. With

the help of Seiberg-Witten map specific method to construct NCSM has been developed [6,7].

To solve the $U(1)$ charge quantization problem, one associates each charge $gq^{(n)}$ of the n th matter field a gauge field $\hat{a}_\mu^{(n)}$ [7]. In the commutative limit, $\theta^{\mu\nu} \rightarrow 0$, $\hat{a}_\mu^{(n)}$ becomes the single gauge field a_μ of the ordinary commuting space-time $U(1)$ gauge theory. But at non-zero orders in $\theta^{\mu\nu}$, $\hat{a}_\mu^{(n)}$ receives corrections and becomes [7]

$$\hat{a}_\xi^{(n)} = a_\xi + \frac{gq^{(n)}}{4}\theta^{\mu\nu}\{a_\nu, \partial_\mu a_\xi\} + \frac{gq^{(n)}}{4}\theta^{\mu\nu}\{f_{\mu\xi}, a_\nu\} + O(\theta^2), \quad (3)$$

where $f_{\mu\nu}$ is the field strength of a_μ . In doing so, the kinetic energy of the gauge boson will, however, be affected. Depending on how the kinetic energy is defined (weight over different field strength of $\hat{a}_\mu^{(n)}$), the resulting kinetic energy will be different even though the proper normalization to obtain the correct kinetic energy in the commutative limit is imposed [7].

In the Standard Model, there are six different matter field multiplets for each generation, i.e. U_R , d_R , $(u, d)_L$, e_R , $(\nu, e)_L$ and (H^0, H^-) , a priori one can choose a different g_i for each of them. After identifying three combinations with the usual g_3 , g_2 and g_1 couplings for the SM gauge groups, there is still freedom to choose different gauge boson self interaction couplings at non-zero orders in $\theta^{\mu\nu}$ (we refer these choices as different schemes). This leads to ambiguities in self interactions of gauge bosons when non-zero order terms in $\theta^{\mu\nu}$ are included. This point was nicely demonstrated in Ref. [7]. It is shown there that with one particular scheme, there are no triple photon self interactions, and with a different scheme there are. This results in non-uniqueness of the theory. This problem needs to be resolved either by performing experiments to test different schemes or applying a underlying theory which uniquely determines the kinetic energy terms. In this paper we propose a solution to this problem from grand unification theory point of view.

The problem with the non-uniqueness of gauge boson kinetic energy is due to the fact that in order to overcome the charge quantization problem, new degrees of freedom have to be introduced. If the degrees of freedom can be reduced and at the same time the correct charge quantization can be obtained, the problem will be solved. To this end we note that

grand unified theory with a single gauge coupling does exactly this. Kinetic energy for SM fields deduced from such a unified theory will be uniquely determined. We will work with $SU(5)$ grand unification [8] to demonstrate this and concentrate on triple gauge boson interactions among the SM fields, photon γ , Z boson and gluons G .

Using Seiberg-Witten map, one can easily obtain the kinetic energy of a noncommutative $SU(5)$ gauge theory. The noncommutative $SU(N)$ gauge boson interactions, when written in terms of commutative conventional gauge fields A_μ^0 and its field strength $F_{\mu\nu}^0 = \partial_\mu A_\nu^0 - \partial_\nu A_\mu^0 - ig_N[A_\mu^0, A_\nu^0]$ to the leading order in $\theta^{\mu\nu}$, are given by [6,9]

$$L = -\frac{1}{2}Tr(F_{\mu\nu}^0 F^{0\mu\nu}) + g_N \theta^{\mu\nu} [\frac{1}{4}Tr(F_{\mu\nu}^0 F_{\rho\sigma}^0 F^{0\rho\sigma}) - Tr(F_{\mu\rho}^0 F_{\nu\sigma}^0 F^{0\rho\sigma})]. \quad (4)$$

The above Lagrangian is uniquely determined to order θ . If a low energy theory is deduced from such a theory, then all the gauge couplings are completely fixed. We now use the noncommutative $SU(5)$ self gauge interactions to derive the NCSM triple neutral gauge boson couplings.

With appropriate Higgs boson representations and corresponding non-zero vacuum expectation values (VEV), such as a 24 and a 5, $SU(5)$ gauge group can be broken down to the SM gauge group, $SU(3)_C \times SU(2)_L \times U(1)_Y$, and then to $SU(3)_C \times U(1)_{em}$. The SM gauge boson self interactions can be obtained by integrating out the heavy degrees of freedom.

The gauge bosons in $SU(5)$ theory are contained in the 24 adjoint representation. In addition to the 8 color gauge bosons G^a , 4 electroweak gauge bosons, $W^\pm$, Z and γ , there are also 12 colored heavy bosons, the X and Y bosons. This representation, in terms of the gauge fields, can be written as [8]

$$\sqrt{2}A_\mu^0 = \sqrt{2}T^a A_\mu^{0a} = \begin{pmatrix} G_\mu^{11} - 2B_\mu/\sqrt{30} & G_\mu^{12} & G_\mu^{13} & \bar{X}_\mu^1 & \bar{Y}_\mu^1 \\ G_\mu^{21} & G_\mu^{22} - 2B_\mu/\sqrt{30} & G_\mu^{23} & \bar{X}_\mu^2 & \bar{Y}_\mu^2 \\ G_\mu^{31} & G_\mu^{32} & G_\mu^{33} - 2B_\mu/\sqrt{30} & \bar{X}_\mu^3 & \bar{Y}_\mu^3 \\ X_\mu^1 & X_\mu^2 & X_\mu^3 & W_\mu^3/\sqrt{2} + 3B_\mu/\sqrt{30} & W_\mu^+ \\ Y_\mu^1 & Y_\mu^2 & Y_\mu^3 & W_\mu^- & -W_\mu^3/\sqrt{2} + 3B_\mu/\sqrt{30} \end{pmatrix}, \quad (5)$$

where T^a are the $SU(5)$ generators with normalization $Tr(T^a T^b) = \delta^{ab}/2$. G_{μ}^{ij} correspond to SM gluon fields, and G^{ii} are linear combinations of the usual G^3 and G^8 gluons. Linear combinations of W_{μ}^3 and B_{μ} give rise to the γ and Z fields. We will indicate the generators of B_{μ} , W_{μ}^3 , and G_{μ}^a fields by T^B , T^{w3} and T^{Ga} , respectively.

To obtain the triple neutral gauge boson self interactions, one needs to expand the $Tr(F^0 F^0 F^0)$ terms in eq. (4). There are only four types of triple neutral gauge boson interactions, BBB , $W^3 W^3 B$, GGB and GGG resulting from $Tr(F^0 F^0 F^0)$ at the order $\theta_{\mu\nu}$. The corresponding traces needed to evaluate the triple neutral gauge boson interactions are

$$\begin{aligned} Tr(T^B T^B T^B) &= \frac{1}{4\sqrt{15}}, & Tr(T^{w3} T^{w3} T^B) &= \frac{3}{4\sqrt{15}}, \\ Tr(T^{Ga} T^{Gb} T^B) &= -\frac{1}{2\sqrt{15}} \delta^{Ga Gb}, & Tr(T^{Ga} T^{Gb} T^{Gc}) &= \frac{1}{4} (if^{GaGbGc} + d^{GaGbGc}), \end{aligned} \quad (6)$$

where f^{abc} and d^{abc} are the totally anti-symmetric and symmetric structure functions for the color gauge $SU(3)_C$ group.

Using the above, we obtain the triple neutral gauge boson interactions as

$$\begin{aligned} L &= \theta^{\mu\nu} g_5 \frac{1}{4\sqrt{15}} \left[\frac{1}{4} B_{\mu\nu} B_{\rho\sigma} - B_{\mu\rho} B_{\nu\sigma} \right] B^{\rho\sigma} \\ &+ \theta^{\mu\nu} g_5 \frac{3}{4\sqrt{15}} \left[\left(\frac{1}{4} W_{\mu\nu}^3 W_{\rho\sigma}^3 - W_{\mu\rho}^3 W_{\nu\sigma}^3 \right) B^{\rho\sigma} \right. \\ &+ \left(\frac{1}{4} W_{\mu\nu}^3 B_{\rho\sigma} - W_{\mu\rho}^3 B_{\nu\sigma} \right) W^{3,\rho\sigma} + \left(\frac{1}{4} B_{\mu\nu} W_{\rho\sigma}^3 - B_{\mu\rho} W_{\nu\sigma}^3 \right) W^{3,\rho\sigma} \left. \right] \\ &- \theta^{\mu\nu} g_5 \frac{1}{2\sqrt{15}} \left[\left(\frac{1}{4} G_{\mu\nu}^a G_{\rho\sigma}^a - G_{\mu\rho}^a G_{\nu\sigma}^a \right) B^{\rho\sigma} \right. \\ &+ \left(\frac{1}{4} G_{\mu\nu}^a B_{\rho\sigma} - G_{\mu\rho}^a B_{\nu\sigma} \right) G^{a,\rho\sigma} + \left(\frac{1}{4} B_{\mu\nu} G_{\rho\sigma}^a - B_{\mu\rho} G_{\nu\sigma}^a \right) G^{a,\rho\sigma} \left. \right] \\ &+ \theta^{\mu\nu} g_5 \frac{1}{4} d^{abc} \left(\frac{1}{4} G_{\mu\nu}^a G_{\rho\sigma}^b G^{c,\rho\sigma} - G_{\mu\rho}^a G_{\nu\sigma}^b G^{c,\rho\sigma} \right). \end{aligned} \quad (7)$$

When the gauge group is further broken down to $SU(3)_C \times U(1)_{em}$ via non-zero VEV of the 5, one obtains the new triple gauge couplings in the physical fields $\gamma(A)$, Z and G . We have,

$$\begin{aligned} L &= \frac{g_5}{4\sqrt{15}} \theta^{\mu\nu} \left\{ c(\theta)(5 - 4c(2\theta)) \left[\frac{1}{4} F_{\mu\nu} F_{\rho\sigma} F^{\rho\sigma} - F_{\mu\rho} F_{\nu\sigma} F^{\rho\sigma} \right] \right. \\ &+ s(\theta)(1 + 4c(2\theta)) \left[\frac{1}{4} (2F_{\mu\nu} Z_{\rho\sigma} F^{\rho\sigma} + Z_{\mu\nu} F_{\rho\sigma} F^{\rho\sigma}) - (2F_{\mu\rho} Z_{\nu\sigma} F^{\rho\sigma} + F_{\mu\rho} F_{\nu\sigma} Z^{\rho\sigma}) \right] \left. \right\} \end{aligned}$$

$$\begin{aligned}
& -c(\theta)(1-4c(2\theta)) \left[\frac{1}{4}(2Z_{\mu\nu}Z_{\rho\sigma}F^{\rho\sigma} + F_{\mu\nu}Z_{\rho\sigma}Z^{\rho\sigma}) - (2Z_{\mu\rho}F_{\nu\sigma}Z^{\rho\sigma} + Z_{\mu\rho}Z_{\nu\sigma}F^{\rho\sigma}) \right] \\
& -s(\theta)(5+4c(2\theta)) \left[\frac{1}{4}Z_{\mu\nu}Z_{\rho\sigma}Z^{\rho\sigma} - Z_{\mu\rho}Z_{\nu\sigma}Z^{\rho\sigma} \right] \Big\} \\
& -\frac{g_5}{8\sqrt{15}}\theta^{\mu\nu} \left[2G_{\mu\nu}^a G_{\rho\sigma}^a (c(\theta)F^{\rho\sigma} - s(\theta)Z^{\rho\sigma}) + (c(\theta)F_{\mu\nu} - s(\theta)Z_{\mu\nu})G_{\rho\sigma}^a G^{a,\rho\sigma} \right. \\
& \left. - 4G_{\mu\rho}^a G_{\nu\sigma}^a (c(\theta)F^{\rho\sigma} - s(\theta)Z^{\rho\sigma}) - 8(c(\theta)F_{\mu\nu} - s(\theta)Z_{\mu\nu})G_{\rho\sigma}^a G^{a,\rho\sigma} \right] \\
& + \frac{g_5}{4}\theta^{\mu\nu} d^{abc} \left(\frac{1}{4}G_{\mu\nu}^a G_{\rho\sigma}^b G^{c,\rho\sigma} - G_{\mu\rho}^a G_{\nu\sigma}^b G^{c,\rho\sigma} \right). \tag{8}
\end{aligned}$$

where $c(n\theta) = \cos(n\theta_W)$, $s(n\theta) = \sin(n\theta_W)$, and $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, $Z_{\mu\nu} = \partial_\mu Z_\nu - \partial_\nu Z_\mu$. Note that each of the coupling and $\sin\theta_W$ given above are at the unification scale. They have to be evolved to the appropriate energy scale depending on the experiment.

It is clear that the triple neutral gauge boson self interactions in the NCSM obtained above are completely specified. The $\gamma\gamma\gamma$, $\gamma\gamma Z$, γZZ , ZZZ , γGG , ZGG and GGG types of interactions are all new compared with the SM predictions. The corresponding couplings are not zero which will lead to observational effects if the energy scale of the noncommutativity is not too high (less than a few TeV) [3].

The triple photon interaction can be studied in various processes, such as $e^+e^- \rightarrow \gamma\gamma$, $e\gamma \rightarrow e\gamma$ and $\gamma\gamma \rightarrow e^+e^-$. The analyses can be carried out in similar ways as those discussed in Ref. [3] with appropriate modifications of the vertices involved. The sensitivity to the noncommutative parameter $\theta^{\mu\nu}$ is typically in the TeV range at next generation of linear colliders with center of mass energy around one TeV [3]. The $\gamma\gamma Z$ coupling can also be studied in the above processes, in $e^+e^- \rightarrow ZZ$, γZ , $Z \rightarrow \gamma\gamma$, and on shell production at $\gamma\gamma \rightarrow Z$ processes. The γZZ coupling can be studied in the processes $e^+e^- \rightarrow ZZ$, γZ . The couplings involving gluons can also be studied at e^+e^- colliders, such as $e^+e^- \rightarrow GG$, and in hadron colliders, such as $pp(\bar{p}) \rightarrow GG$, and in $Z \rightarrow GG$ decay. All the processes having gluons in the final state may be more difficult to study compared with the ones without due to hadronization. But if sophisticated polarization analyses can be carried out, these processes can also provide interesting information about the noncommutativity of space-time.

In conclusion, in this paper we have proposed a solution to the arbitrariness of kinetic energy problem of the noncommutative Standard Model from grand unification point of view. The kinetic energy and the resulting self gauge boson interactions are uniquely determined in noncommutative $SU(5)$ theories. Although we have only worked with $SU(5)$ grand unification as an example, the conclusion will hold in any grand unification theories with a single coupling constant. At the first order in the noncommutative parameter $\theta^{\mu\nu}$, there are new triple gauge boson interactions, such as $\gamma\gamma\gamma$, $\gamma\gamma Z$, γZZ , ZZZ , γGG , ZGG and GGG . These interaction are forbidden in the Standard Model. The theory can be tested by experiments. Non observation of these interactions can also put bounds on the noncommutative parameter $\theta^{\mu\nu}$.

We thank P.-M. Ho for discussions. This work was supported in part by DOE grant DE-FG03-96ER40969 and by National Science Council under grants NSC 89-2112-M-002-058 and NSC 89-2112-M-002-065, and in part by the Ministry of Education Academic Excellence Project 89-N-FA01-1-4-3. One of us (NGD) would like to thank hospitality provided by the Department of Physics at National Taiwan University where this work was done.

REFERENCES

- [1] H.Snyder, Phys. Rev. **71**, 38(1947); ibid **72**, 68(1947; A. Connes, Noncommutative Geometry, Academic Press (1994); M. Douglas and N. Nekrasov, hep-th/0106048.
- [2] I. Riad and M. Sheikh-Jabari, JHEP, **0008**, 045(2000).
- [3] J. Hewett, F. Petriello and T. Rizzo, hep-ph/0010354, Phys. Rev. **D64**, 075012(2001); P. Mathews, hep-ph/0011332, Phys. Rev. **D63**, 075007(2001); S. Baek et al., hep-ph/0103068, Phys. Rev. **D64**, 056001(2001).
- [4] M. Hayakawa, Phys. Lett. **B478**, 394(2000); K. Matsubara, Phys. Lett. **B482**, 417(2000).
- [5] J. Seiberg and E. Witten, JHEP, **9909**, 032(1999).
- [6] J. Madore et al., hep-th/0001203, Eur. Phys. J. **C16**, 161(2000); B. Jurco et al., hep-th/0006246, Eur. Phys. J. **C17**, 521(2000); B. Jurco. P. Schupp and J. Wess, hep-th/0102129, Nucl. Phys. **B604**, 148(2001); B. Jurco et al., hep-th/0104153, Eur. Phys. J. **C21**, 383(2001).
- [7] X. Calmet et al., hep-ph/0111115.
- [8] H. Georgi and S. Glashow, Phys. Rev. Lett. **32**, 438(1974).
- [9] C. Carlson, C. Carone and R. Lebed, hep-ph/0107291, Phys. Lett. **B518**, 201(2001).